




# *Results from ALICE*

*Christine Nattrass  
for the ALICE collaboration  
University of Tennessee at Knoxville*







# Exploring QCD at high temperatures



Quark-gluon plasma




Deconfined - more degrees of freedom



Confined - fewer degrees of freedom

**Size:** 16 x 26 meters  
**Weight:** 10,000 tons  
**Detectors:** 18






# Bulk properties



# Bulk properties

Collision system on the slide




pp



# Bulk properties

Collision system on the slide




**Pb+Pb**




# $dN_{ch}/d\eta$ versus $\sqrt{s}$

$dN_{ch}/d\eta$  versus  $\sqrt{s}$



Relative increase in  $dN_{ch}/d\eta$




## Results:

- increase with energy significantly stronger in data than MC's
- ALICE & CMS agree to within  $1 \sigma$  ( $< 3\%$ )

Eur. Phys. J. C (2010) 68: 345–354  
 Eur. Phys. J. C (2010) 68: 89–108  
 Eur. Phys. J. C (2010) 65: 111-125




# Centrality dependence of $dN_{ch}/d\eta$







# Centrality dependence of $dN_{ch}/d\eta$




RHIC data  
scaled by 2.1

PHENIX  
PRC 71, 034908 (2005)

Pb+Pb 9



# Transverse Energy



PRC71:034908 (2005)  
PRC70:054907 (2004)

Centrality dependence similar to  
RHIC (PHENIX)

- $E_T^{\text{had}}$  from charged hadrons directly measured by the tracking detectors
- $f_{\text{total}}$  from MC to convert into total  $E_T$
- From RHIC to LHC
  - $\sim 2.5$  increase  
 $dE_T/d\eta / (0.5 * N_{\text{part}})$
- Energy density (Bjorken)

$$\varepsilon = \frac{1}{\pi R^2 \tau} \frac{dE_t}{dy} \quad R = 1.12 A^{1/3} \text{ fm}$$



- $\varepsilon \tau \sim 16 \text{ GeV}/(\text{fm}^2 \text{c})$   
RHIC:  $\varepsilon \tau = 5.4 \pm 0.6 \text{ GeV}/(\text{fm}^2 \text{c})$



# $\sqrt{s_{NN}}$ dependence

- $dN_{ch}/d\eta/(0.5*N_{part}) \sim 8$
- **2.1 x RHIC**  
1.9 x pp (NSD) at 2.36 TeV
- growth with  $\sqrt{s}$  faster in AA than pp
- $dE_T/d\eta/(0.5*N_{part}) \sim 9$  in 0-5%
- $\sim 5\%$  increase of  $N_{part}$  (353  $\rightarrow$  383)  
 $\rightarrow$  **2.7 x RHIC**  
(consistent with 20% increase of  $\langle p_T \rangle$ )

Grows faster than simple logarithmic scaling extrapolated from lower energy




**Pb+Pb** 11




$$E_T / N_{ch}$$

- Consistent behavior for  $E_T$  and  $N_{ch}$



- Both increase with energy
- Both show steady rise from peripheral to central
- $E_T / N_{ch}$  independent of centrality
- $E_T / N_{ch}$  slightly increases with energy





Pb+Pb<sup>12</sup>



# Charged particle spectra




# Charged particle spectra





# Nuclear modification factor ( $R_{AA}$ )




$$R_{AA}(p_T) = \frac{(1/N_{evt}^{AA}) d^2 N_{ch}^{AA} / d\eta dp_T}{\langle N_{coll} \rangle (1/N_{evt}^{PP}) d^2 N_{ch}^{PP} / d\eta dp_T}$$

**Pb+Pb** 15



# Nuclear modification factor ( $R_{AA}$ )






$$R_{AA}(p_T) = \frac{(1/N_{evt}^{AA}) d^2 N_{ch}^{AA} / d\eta dp_T}{\langle N_{coll} \rangle (1/N_{evt}^{pp}) d^2 N_{ch}^{pp} / d\eta dp_T}$$

**Pb+Pb** 16







# Particle identification







# Identified particle spectra






# $\bar{p}/p$ ratio in p+p collisions



0.9 TeV:  $\bar{p}/p = 0.957 \pm 0.006(\text{stat}) \pm 0.014(\text{syst})$   
 7 TeV:  $\bar{p}/p = 0.990 \pm 0.006(\text{stat}) \pm 0.014(\text{syst})$




# Identified Particle spectra in Pb-Pb collisions






# Identified Particle spectra in Pb-Pb collisions





# Identified Particle spectra in Pb-Pb collisions









# Strange particle spectra



# Strange particles in pp collisions




- **PYTHIA and PHOJET consistently below data**






# Stranger particles



ALI-PREL-337




# Particle ratios in pp collisions



Thermus fit fails – worked better at lower energies




# Nuclear modification factor ( $R_{AA}$ )



$$R_{AA}(p_T) = \frac{(1/N_{evt}^{AA}) d^2 N_{ch}^{AA} / d\eta dp_T}{\langle N_{coll} \rangle (1/N_{evt}^{PP}) d^2 N_{ch}^{PP} / d\eta dp_T}$$



# Baryon anomaly: $\Lambda/K_S^0$






# Heavy flavor spectra




# Non-photonic electrons

## Beauty + Charm




## Beauty Electrons






# Charm cross section





# Charm nuclear modification factor








# J/Ψ nuclear modification factor

Inclusive J/ψ  $R_{AA}^{0-80\%} = 0.49 \pm 0.03$  (stat.)  $\pm 0.08$  (sys.)



ALI-PREL-3779



# Conclusions

- Charged particle multiplicities and transverse energy
  - pp: higher than model predictions
  - Pb-Pb: higher than model predictions, centrality dependence similar to RHIC
- Charged particle spectra
  - pp: excellent PID measurements to low  $p_T$ , measurements of  $\bar{p}/p$  ratio, failure of statistical models
  - Pb+Pb: suppression greater than RHIC, comparable suppression to RHIC at same  $dN_{ch}/d\eta$ , failure of hydro models to describe protons
- Strange particles
  - pp: models fail significantly
  - Pb+Pb: baryon enhancement,  $\Lambda$  &  $K_S^0$  suppression similar
- Heavy flavor
  - pp: charm cross sections measured, separation of charm & beauty
  - Pb+Pb: suppression of heavy flavor similar to charged particles



# Many results not covered

---


- HBT correlations
- Hydrodynamical flow
- Di-hadron correlations
- Charged track jets
- Neutral mesons
- Resonances
- Diffraction in pp
- Ultraperipheral Pb+Pb collisions
- CP violation
- $p_T$  fluctuations



# Backup slides

---



# p+p collisions





# The ALICE Collaboration

- ~1000 Members  
63% from CERN  
member states
- ~30 Countries
- ~100 Institutes
- ~150 MCHF capital cost  
(+magnet)



## ***US ALICE***

11 Institutions 53 members (inc. 12 grad. Students)  
*Cal. St. U. –San Luis Obispo, Creighton University, University of Houston, Lawrence Berkeley Nat. Lab, Lawrence Livermore Nat. Lab, Oak Ridge Nat. Lab, Ohio State University, Purdue University, University of Tennessee, Wayne State University, Yale University*







# EMCal

Current coverage:

$\Delta\eta=1.4, \Delta\phi=39^\circ$  ( $R \approx 0.3$  max)

Full calorimeter installation scheduled for 2012



- Lead-scintillator sampling calorimeter
- 13 k towers
- Each tower  $\Delta\eta \times \Delta\phi = 0.014 \times 0.014$
- Shashlik geometry
- Avalanche photodiodes
- $\Delta\eta=1.4, \Delta\phi=107^\circ$
- $\sigma(E)/E = 0.12/\sqrt{E} + 0.02$





# ALICE detectors and acceptance

## Central barrel- $0.9 < \eta < 0.9$

- $\Delta\phi = 2\pi$  tracking, PID (TPC/ITS/TRD/ToF)
- single arm RICH (HMPID)
- single arm e.m. cal (PHOS)
- jet calorimeter (EMCal)

## Forward muon arm- $2.4 < \eta < -4$



- absorber, 3 T-m dipole magnet
- 5 tracking + 2 trigger planes

## Multiplicity detectors- $3.4 < \eta < 5$

- including photon counting in PMD


## Trigger & timing detectors

- 6 Zero Degree Calorimeters
- T0: ring of quartz window PMT's
- V0: ring of scintillator Paddles






# Charged Particle $R_{AA}$




$$R_{AA}(p_T) = \frac{(1/N_{evt}^{AA}) d^2 N_{ch}^{AA} / d\eta dp_T}{\langle N_{coll} \rangle (1/N_{evt}^{pp}) d^2 N_{ch}^{pp} / d\eta dp_T}$$

# *Elliptic flow at 2.76 TeV*

Phys. Rev. Lett. 105, 252302 (2010)



# *Elliptic flow at 2.76 TeV*






# $dN_{ch}/d\eta$ in Pb-Pb collisions at $\sqrt{s}_{NN}=2.76$ TeV

Measured  $dN_{ch}/d\eta = 1584 \pm 76$  (sys.)

PRL, 105, 252301 (2010)

Post-pp



Monte Carlo, coherence via collectivity, strong gluon saturation

Saturation ideas

Data driven, limiting frag.

Miscellaneous: superposition, WNM, diffusion eqs., DPM + shadowing/percolation

Compilation from N. Armesto

Pb+Pb 45